

# Does Dark Chocolate Relieve Menstrual Pain in Adult Women?: A Study Among Indian Population

Amita Verma<sup>1</sup>, Archana Kadam<sup>2</sup>

<sup>1</sup>Assistant Professor, <sup>2</sup>Associate Professor, Dept. of Physiology, Dr. D.Y. Patil Medical College, Navi Mumbai

## Abstract

**Introduction:** Premenstrual pain (PMS) and Dysmenorrhea or painful menstruation is defined as a severe, painful, cramping sensation in the lower abdomen that is often accompanied by other symptoms, such as sweating, headaches, nausea, etc all occurring just before or during the menses<sup>1</sup>. A variety of drugs, predominantly non steroidal anti-inflammatory drugs (NSAIDS) are used to provide analgesia during the period. The present study was undertaken to evaluate the claims of Dark chocolate as an alternative to NSAIDS.

**Methodology:** 90 selected students were divided into 3 groups and were provided 120gms/day of dark chocolate, milk chocolate and no chocolate based on their group. Numeric Rate Scale (NRS) was used to measure pain before and after intervention.

**Result:** Pre menstrual pain and menstrual pain was significantly reduced after the consumption of Dark chocolate.

**Keywords:** Premenstrual pain, non steroidal anti-inflammatory drugs, Numeric Rate Scale.

## Introduction

Adolescence in girls has been recognized as a special period which signifies the transition from girlhood to womanhood and is marked with the onset of menarche, an important milestone which is often associated with problems of irregular menstruation, excessive bleeding and dysmenorrhoea.<sup>3,4</sup> Dysmenorrhea or painful menstruation is defined as a severe, painful, cramping sensation in the lower abdomen that is often accompanied by other symptoms, such as sweating, headaches, nausea, vomiting, diarrhoea, and tremulousness, all occurring just before or during the menses.<sup>1</sup> Dysmenorrhea may be categorized into two types as primary and secondary. Primary dysmenorrhea is defined as painful menses among females with normal pelvic anatomy, frequently beginning during adolescence. It is observed only in ovulatory cycles, frequently emerging within 6 to 12 months after menarche with no pathology or organic

basis. Secondary dysmenorrhea is a menstrual pain associated with underlying pathology and its onset might be years after menarche.<sup>5</sup> The cause of primary dysmenorrhea is not well established. However, the responsible cause has been identified on the hyper-production of uterine prostaglandins, particularly of PGF 2 $\alpha$  and PGE 2, thus resulting in increased uterine tone and high-amplitude contractions.<sup>6</sup>

Women with dysmenorrhea have higher levels of prostaglandins, during the first two days of menses.<sup>7</sup> under the control by progesterone while immediately prior to menstruation, prostaglandin levels increase.<sup>6, 8</sup>

The levels of prostaglandin F2 $\alpha$  are especially high during the first two days of menstruation in women with severe primary dysmenorrhea<sup>9</sup> who are treated by use of NSAIDs such as aspirin, ibuprofen and naproxen, or by use of Prostaglandin inhibitors which though being better in pain relief cause GI disturbances and have a propensity for severe side effects on long term use.<sup>1, 10, 11</sup>

## Corresponding Author:

**Archana Kadam**

Associate Professor, Dept. of Physiology,  
Dr. D.Y. Patil Medical College, Navi Mumbai  
e-mail: drarchanashirsath24@gmail.com

There is some evidence supporting the role of some dietary supplements, including Omega-3 fatty acids, such as fish oil, vitamin B1, vitamin B6, vitamin D, vitamin E, magnesium and vitamin B6, on menstrual

pain relief.<sup>12, 13, 14</sup> Disruption of calcium regulation has been proposed as an underlying factor for increasing incidence and severity of menstrual pain<sup>15, 16</sup>

Chocolate is rich in vitamin A, B1, C, D, and vitamin E. In addition, chocolate also contains antioxidants of phenol and flavonoids, rich in minerals such as calcium, potassium, iron, a little omega 3 and 6, and high magnesium<sup>17</sup> which can reduce menstrual pain and premenstrual occurrence in women. Because of the mentioned benefits of dark chocolate, this study was undertaken to examine the effect of dark chocolate on menstrual pain in adolescent women.

## Materials & Method

**Study Design:** This was experimental study with randomized testing with control group design.

The study was conducted on medical students at DY Patil Medical College Navi Mumbai. 90 samples were recruited by random sampling technique. The students who had given the history of severe primary dysmenorrhea were selected for the present study.

### Inclusion criteria:

- Single, Young woman aged 18-21 years
- Experiencing severe pre menstrual and menstrual pain in the last 6months
- Regular menstrual cycle
- Not taking analgesic medication in last 24 hours

**Exclusion criteria:** Students with history of any chronic illnesses, or those using any contraceptive pills or any vitamin supplements

**Intervention:** 90 students that were selected were divided into 3 groups of 30 students each.

**Group 1:** Received 120gms/day of dark chocolate for 3 days.

**Group 2:** Received 120gms/day of milk chocolate for 3days

**Group 3:** With severe dysmenorrhea who did not receive any chocolate.

Doses given from onset of PMS till Second day of menses.

**Instrument:** Numeric Rate Scale (NRS)<sup>2</sup> was used to measure pain. The 10-point numeric scale ranges

from '0' representing one pain extreme (e. g. "no pain") to '10' representing the other pain extreme Pain was measured before and after intervention.

Institutional ethical committee clearance was obtained prior to commencement of the study. No data of identification apart from weight, height were taken on the subject information sheet. The data of intervention pain scales were tabulated on MS Excel data sheet and compiled. The statistical analysis was done using SPSS software version 16.




Fig 01: Pain Scale Index

## Observations

The present study comprised of 90 subjects, females of 18 to 21 years of age, with the mean age being 19 years and 3 months. No statistically significant differences were found in the age and BMI of the selected subjects. The group I consisted of individuals who were given dark chocolate during their study period. The details of group I are as follows (Table 1)

Table 1: Dark Chocolate Group (Group I)

Subjects	Mean Age (Years)	Mean PSI (Pre)	Mean PSI (Post)	Mean BMI
30	19.3	8.36	7.02	23.05

The subjects showed a improvement in their pain scales with dark chocolate. Paired T Test (two tailed) was conducted to ascertain the significance, which revealed P value is less than 0.0001. By conventional criteria, this difference is considered to be extremely statistically significant at 95% confidence interval.

Group II comprised of 30 subjects, who were administered milk chocolate as a part of intervention to assess its effects on pain scale index during menstrual pain. (table 02)

Table 2: Milk Chocolate Group (Group II)


Subjects	Mean Age (Years)	Mean PSI (Pre)	Mean PSI (Post)	Mean BMI
30	19.6	8.41	7.37	22.52

In terms of pain, the subjects showed a mild improvement with administration of milk chocolate. Paired T Test (two tailed) was conducted to ascertain the significance, which revealed that the P value < 0.0001. By conventional criteria, this difference is considered to be statistically significant at a confidence interval of 95%.

**Table:3 Control Group (Group I)**

Subjects	Mean Age (Years)	Mean PSI	Mean BMI
30	19.3	8.43	22.71

### Mean Pain Scale Levels



One way ANOVA analysis was conducted to ascertain the variance among the three groups which revealed that the f-ratio value is 5.15115. The p-value is 0.007687. The result is significant at 95 % confidence interval indicating that a variance exists between all three groups with dark chocolate showing highest mean difference in reduction of pain levels (1.34) as compared to other two groups.

In Summary, the study revealed that pain levels among adolescent females experiencing dysmenorrhea was reduced on consumption of dark chocolate, in comparison with milk chocolate and no chocolate.

## Discussion

This study was conducted to compare the effects of administration of dark chocolate and milk chocolate on primary dysmenorrhea. The results showed that obtained efficacy in pain reduction was higher with taking dark chocolate and moderate with taking milk chocolate.

Enhanced release of PGs, allegedly from disintegrating cells during endometrial sloughing, is believed to cause myometrial hyper contractility, resulting in ischemia and hypoxia of the uterine muscle, and, ultimately, pain.<sup>18, 19</sup>

All women have increased levels of PGs during the luteal phase compared with the follicular phase of

ovulatory cycles. However, compared with eumenorrheic women, dysmenorrheic women have higher levels of PGs, as measured in luteal phase endometrial biopsies, jet washings and menstrual fluids.<sup>20, 21</sup> Higher circulating levels of PGs (PGF2 $\alpha$  and PGE2) have been reported in women with dysmenorrhea compared with asymptomatic women during menstruation, and these PG levels are highest during the first 48 h of menses, when symptoms peak.<sup>18, 22, 23, 24</sup> Further, the severity of menstrual pain and associated symptoms of dysmenorrhea are directly proportional to the amount of PGs released.<sup>21, 25</sup> In addition, clinical administration of exogenous PGs results in uterine contraction and often also produces the same systemic symptoms that frequently accompany dysmenorrhea.<sup>24, 26</sup>

Magnesium influences the contractility, tone and relaxation of the uterine smooth muscle; and may inhibit the synthesis of prostaglandin based on inhibition of biosynthesis of prostaglandin. Studies have revealed that menstrual pain is relieved by good diet.<sup>27, 28, 29, 30</sup> Paath, (2004) stated adolescent women need to maintain a good nutritional status, with a balanced way of eating at the time of menstruation.<sup>31</sup>

It has been reported that plasma levels of calcium in premenstrual period is lower in people suffering from premenstrual syndrome.<sup>32, 33</sup> Di Cintio et al reported a relationship between calcium intake in food and reducing the severity of primary dysmenorrhea. Also, a small randomized trial showed that increasing dietary calcium intake reduce mood and pain symptoms associated with menstruation<sup>34, 35</sup>

Repeated monthly painful episodes may lead to the development of central sensitivity to pain.<sup>36, 37</sup> Central sensitization is defined as an abnormal augmentation of pain by mechanisms within the central nervous system (CNS), and therefore represents a state where the response to normal peripheral inputs is greatly enhanced.<sup>38, 39</sup> Primary dysmenorrhea has been classified as a member of the central sensitivity syndromes together with several other clinical conditions including fibromyalgia and tension-type headaches.<sup>36, 37</sup> These syndromes are characterized by pain hypersensitivity in the absence of tissue injury, inflammation, or a lesion to the nervous system.<sup>36, 39</sup>

In a previous study examining the benefits of dark chocolate to reduce menstrual pain stated that the chocolate content of GABA (Gamma-Amino Butyric acid) is an amino acid that has

a major function as a neurotransmitter in the central nervous system. Gamma aminobutyric acid inhibits nerve transmission in the brain, calming nervous activity. The 5-HT creates a sense of comfort and increase serotonin levels. L-tyrosine gives rise to body-wide relaxation. L-glutamine can be utilized directly by the brain for energy production in brain cells. Thus it can improve brain function and facilitate the utilization of GABA and S-HTP. In theory, magnesium has a direct effect on vascular pressure and can regulate the entry of calcium into the smooth muscle cells of the uterus, so magnesium affects contraction and relaxation of smooth muscle of the uterus.<sup>41</sup>

Magnesium can also suppress inflammation by inhibiting the formation of prostaglandins. So in the group treated with dark chocolate states menstrual pain is reduced significantly.<sup>41</sup>

Copper is a cofactor for a number of enzymes and is required for iron transport and other process.<sup>43, 44</sup> Dark chocolate provides 31% whereas milk chocolate provides 10% of the U. S. RDA for copper per 100-kcal serving, dark chocolate provides 25% of the RDA (1.90 mg) and Milk chocolate contains 5% of the RDA for iron.<sup>42</sup> A study in 2005 stated, dark chocolate contains complex carbohydrates, antioxidants (flavonoid polyphenols), vitamin B6, unsaturated fatty acids (omega 3 and omega 6) and minerals (magnesium, calcium, iron) that influence and regulate menstrual cycle by balancing the levels of the sex hormones in the blood during the luteal phase of the cycle.<sup>44</sup>

**Conclusion:** The study concluded that in the selected study group, the pain levels during dysmenorrhoea was significantly reduced by administration of dark chocolate as compared to no administration. The use of a natural substitute is a viable option to NSAIDS as it carries none of the side effects associated with the drugs. The study is limited by the small sample size and lack of variability in the age group of the subjects. A wider study is needed to effectively apply this in the entire population.

**Conflict of Interest:** None

**Source of Funding:** Self Funded

## References

1. Lobo RA, Gershenson DM, Lentz GM, Valea FA. Comprehensive Gynecology E-Book. Elsevier Health Sciences; 2016 Jun 22.
2. Judha M, Sudarti FA. Teori pengukuran nyeri dan nyeri persalinan. Jogjakarta, Muha Medika. 2012.
3. Agarwal AK, Agarwal A. A study of dysmenorrhea during menstruation in adolescent girls. Indian journal of community medicine: official publication of Indian Association of Preventive & Social Medicine. 2010 Jan; 35 (1):159.
4. Dhingra R, Kumar A, Kour M. Knowledge and practices related to menstruation among tribal (Gujjar) adolescent girls. Studies on Ethno-Medicine. 2009 Jan 1; 3 (1):43-8.
5. Sharma N, Sagayaraj M, Sujatha B. Menstrual characteristics and prevalence of dysmenorrhea in college students. International Journal of Scientific and Research Publications. 2014; 4 (10):1-6.
6. Iacovides S, Avidon I, Baker FC. What we know about primary dysmenorrhea today: a critical review. Human reproduction update. 2015 Sep 7; 21 (6):762-78.
7. Dawood MY. Primary dysmenorrhea: advances in pathogenesis and management. Obstetrics & Gynecology. 2006 Aug 1; 108 (2):428-41.
8. Wong CL, Farquhar C, Roberts H, Proctor M. Oral contraceptive pill as treatment for primary dysmenorrhoea. Cochrane Database of Systematic Reviews. 2009 (2).
9. Nair P, Grover VL, Kannan AT. Awareness and practices of menstruation and pubertal changes amongst unmarried female adolescents in a rural area of East Delhi. Indian journal of community medicine. 2007 Apr 1; 32 (2):156.
10. Morrison BW, Daniels SE, Kotey P, Cantu N, Seidenberg B. Rofecoxib, a specific cyclooxygenase-2 inhibitor, in primary dysmenorrhea: a randomized controlled trial. Obstetrics & Gynecology. 1999 Oct 1; 94 (4):504-8.
11. Smith WL, Langenbach R. Why there are two cyclooxygenase isozymes. The Journal of clinical investigation. 2001 Jun 15; 107 (12):1491-5.
12. Liu CZ, Xie JP, Wang LP, Zheng YY, Ma ZB, Yang H, Chen X, Shi GX, Li SL, Zhao JP, Han JX. Immediate analgesia effect of single point acupuncture in primary dysmenorrhea: a randomized controlled trial. Pain Medicine. 2011 Feb 1; 12 (2):300-7.
13. Penland JG, Johnson PE. Dietary calcium and manganese effects on menstrual cycle symptoms. American journal of obstetrics and gynecology. 1993 May 1; 168 (5):1417-23.



14. Charandabi SM, Mirghafourvand M, Chegini SN, Javadzadeh Y. Calcium With and Without Magnesium for Primary Dysmenorrhea: A Double-Blind Randomized Placebo Controlled Trial. *Int J Women's Health Reprod Sci.* 2017 Oct 1; 5 (4):332-8.
15. Alvir JM, Thys-Jacobs S. Premenstrual and menstrual symptom clusters and response to calcium treatment. *Psychopharmacology bulletin.* 1991.
16. Thys-Jacobs S. Micronutrients and the premenstrual syndrome: the case for calcium. *Journal of the American College of Nutrition.* 2000 Apr 1; 19 (2):220-7.
17. Nurazizah E, Tih F, Suwindere W. Black chocolate consumption reduces subjective symptoms in 18-22 years old females with premenstrual syndrome. *Journal of Medicine and Health.* 2015 Feb 27; 1 (1).
18. Dawood MY. Dysmenorrhea and prostaglandins. *In Gynecologic endocrinology* 1987 (pp. 405-421). Springer, Boston, MA.
19. Hayaishi O, Matsumura H. Prostaglandins and sleep. *Adv Neuroimmunol* 1995; 5:211-216.
20. Chan WY, Hill JC. Determination of menstrual prostaglandin levels in nondysmenorrheic and dysmenorrheic subjects. *Prostaglandins* 1978; 15:365-375.
21. Lundstrom V, Green K. Endogenous levels of prostaglandin F2alpha and its main metabolites in plasma and endometrium of normal and dysmenorrheic women. *Am J Obstet Gynecol* 1978; 130:640-646.
22. Hofmeyr GJ. Dysmenorrhoea. In: Bassin J (ed). *Topics in Obstetrics and Gynaecology.* Johannesburg: Julmar Communications, 1996, 269-274.
23. Coco AS. Primary dysmenorrhea. *Am Fam Physician* 1999; 60:489-496.
24. Chan WY, Dawood MY, Fuchs F. Prostaglandins in primary dysmenorrhea. Comparison of prophylactic and nonprophylactic treatment with ibuprofen and use of oral contraceptives. *Am J Med* 1981; 70:535-541.
25. Dawood MY. Dysmenorrhea. *Endometrium* 1995; 6:363-377.
26. Lavin, N., *Manual of Endocrinology and Metabolism.* Boston: Little, Brown; (1986). Website available at:<http://www.franco-lania.com/manual-of-endocrinology-and-metabolism-lavin.pdf>
27. Al-Ghamid, S. M., Cameron, E. C., and Sutton, RA., Magnesium deficiency: pathophysiologic and clinical overview. *Am J Kidney Dis* (1994) 24:737-52
28. Walker, S.M., In vitro synthesis of prostaglandin F2a by human endometrium. Abstracts of the Twenty-first British Congress of Obstetrics and Gynecology, (1977) P 37. Sheffield. Website available at:<http://iasj.net/iasj?func=fulltext&aId=72026>
29. Andrea J, Candace N. Pelvic Pain and Dysmenorrhea. In: Novak E, Berek JS. *Berek & Novak's Gynecology.* 14th ed. Philadelphia: Lippincott Williams & Wilkins; 2007:506-534.
30. Paath, E. F. (2004) *Gizi dalam Kesehatan Reproduksi.* Jakarta: EGC
31. Muneyvirci-Delale O, Nacharaju VL, Altura BM, Alutra BT. Sex steroid hormones modulate serum ionized magnesium and calcium levels throughout the menstrual cycle in women. *Fertil Steril.* 1998; 69 (5):958-962.
32. Thys-Jacobs S, McMahon D, Bilezikian JP. Cyclical changes in calcium metabolism across the menstrual cycle in women with premenstrual dysphoric disorder. *J Clin Endocrinol Metab.* 2007; 92 (8):2952-2959
33. Di Cintio E, Parazzini F, Tozzi L, et al. Dietary habits, reproductive and menstrual factors and risk of dysmenorrhoea. *Eur J Epidemiol.* 1997; 13 (8):925-930.
34. Penland JG, Johnson PE. Dietary calcium and manganese effects on menstrual cycle symptoms. *Am J Obstet Gynecol.* 1993; 168 (5):1417-1423.
35. Yunus MB. Fibromyalgia and overlapping disorders: the unifying concept of central sensitivity syndromes. *Semin Arthritis Rheum* 2007; 36:339-356.
36. Yunus MB. Central sensitivity syndromes: a new paradigm and group nosology for fibromyalgia and overlapping conditions, and the related issue of disease versus illness. *Semin Arthritis Rheum* 2008; 37:339-352.
37. Woolf CJ. Pain: moving from symptom control toward mechanism-specific pharmacologic management. *Ann Intern Med* 2004; 140:441-451

38. Woolf CJ. Central sensitization: uncovering the relation between pain and plasticity. *Anesthesiol* 2007; 106:864-867.
39. Sessle BJ. What is pain, and why and how do we experience pain? In: Lavigne G, SessleBJ, Choiniere M, Soja PJ (eds). *Sleep and Pain*. Seattle: IASP Press, 2007, 23-43.
40. Smith, G. Z., & Smith, C. C. (2012). *Chocolate candies fortified with natural amino acids and/or herbal nutrients for relief of insomnia, pms, and difficulty concentrating: Google Patents*
41. National Academy of Sciences. Institute of Medicine. *Dietary Reference Intakes (DRIs): Recommended Intakes for Individuals, Vitamins*. 2004184. National Academy of Sciences. Institute of Medicine. *Dietary Reference Intakes (DRIs): Recommended Intakes for Individuals, Vitamins*. 2004
42. Olivares M. Uauy R. Copper as an essential nutrient. *Am J Clin Nutr*. 1996; 63:791S-796S.
43. Uauy R. Olivares M. Gonzalez M. Essentiality of copper in humans. *Am J Clin Nutr*. 1998; 67:952S-959S.
44. Speroff, L., & Fritz, M. A. (2005). *Clinical gynecologicendocrinology and infertility*: lippincottWilliams & wilkins